



DETERMINATION OF MEDIAN TOLERANCE LIMIT (LC₅₀) OF CLARIAS BATRACHUS FOR MERCURIC CHLORIDE

Bhise J.V.

Dr.Manorama & Prof. H.S Pundkar Arts, Comm. & Science College,Balapur,Dist.Akola

Abstract

Industries are the major sources of heavy metal pollution and it is released into water and soil. Heavy metals cause several ill effects to aquatic living organisms and environment. The acute toxicity of Mercury against fresh water *Clarias batrachus* and the chronic toxicity of sublethal concentrations of heavy metals on the different organs such as liver, gill, muscle, kidney and brain of the chosen fresh water fish. Adult cat fish treated separately with Mercury Chloride (HgCl₂) for 24, 48, 72, and 96hr at different dose levels showed significant behavioral changes in the Lethal concentration were determined by probit analysis method. The LC₅₀ values for HgCl₂ at 96 hours was found to be 1.85ppm; respectively. Hence the toxicants selected, HgCl₂ is highly toxic in cat fish.

Keywords: *Clarias batrachus*, Lethal concentration, Mercuric chloride

Introduction

In India, there has been a tremendous space in the development and agricultural practices since independence with the progress in industrial development due to advancement of technology; environmental pollution has grown in alarming proportion. The problem of pollution of the water where the wastes are usually discharged has increased to a great extent in recent years. Aquatic life is strongly influenced by physical properties of a water body. It is known that heavy metals as well as agro-pollutants are potentially harmful to the aquatic lives. All pesticides applied for the pest control eventually pollute the water resources either in their original chemical form or in some degraded variety. On the other hand, all industries discharge their effluents indiscriminately in the adjoining water areas

and frequently cause serious hazards to aquatic life. Among the aqua fauna, fishes are affected to a significant extent. There are ample reports dealing with mortality and pollution [1].

Short term toxicity tests had been designed for the estimation of lethality of the xenobiotics to the test specimens. The most widely adapted technique for short term bio assay is the evaluation of a Medium Tolerance Limit (TLM). Determination of 24, 48, 72 and 96h TLM value by probit analysis (static bio assay) procedure is generally applied to determine the toxicity levels of the xenophobic materials to various aquatic organisms under laboratory conditions [2]. These toxicity tests are necessary to predict the safe concentration of the chemicals in the environment [3]. TLM or Lethal Concentration₅₀ (LC₅₀) is the particular concentration of a chemical in the experimental water at which just 50% of the test organisms are able to survive for a specific period of exposure [4],[5]. Many investigators had accepted this method for the determination of acute toxicity[6], [7], [8] even in recent years for scientific, economic and ethical reasons the TLM test have been prescribed as one of the toxicity tests. APHA [4] suggested that the preferable duration for a toxicity test should be at least 24, 48, 72 and 96h. For the present study, the selected exposure period was 96 in the fish *Clarias batrachus* [9].

Materials and Methods

Clarias batrachus obtained from local fishermen were weighing about 50±5 g and were acclimatized in the laboratory condition for 2 weeks. Fishes were feed with fish Food. Mercuricchloride (HgCl₂) was chosen as xenobiotics for estimation of their lethality on *C. batrachus*. A batch of 10 adults, healthy disease - free laboratory acclimated individual

of some age irrespective of sex, were released in plastic aquaria measuring 10 L water. The test fish were starved for 24, 48, 72, and 96h; Test period.

Feeding tends to increase the metabolic activities such as rate of respiration, excretion and production of other waste products altogether affect the toxicity of the test solution, so food supply to the test specimen was avoided. For the determination of LC50 at 96h for HgCl₂ a set of 4 plastic aquaria were arranged for each chemical, containing 10 of water. After releasing the test individuals, different concentrations of selected chemical were added to respective aquaria very carefully and mixed thoroughly.

Any change in behavior and other responses were carefully noted. The specific doses had been determined by trial and error method. The individuals were considered dead when they failed to respond to touch stimulus. After the death of each fish, its body was removed immediately from the containers along with water allotted for each fish. The aquaria water containing the compounds was renewed after each 24h keeping the concentration constant throughout the experiment. After 96h (the end of the test), survived individuals in each container were counted. Determination of LC50 at 96h values of HgCl₂, in adult *C. batrachus* had been done by the Finney [8] method of probity analysis [10].


Result and Discussion

Fish exposed to different doses of pollutants displayed marked behavioral changes. Fish exposed to sub lethal doses of the test chemicals exhibit cessation of all movements and became very sluggish. Toxicity is a relative property of a chemical which refers to its potency to induce harmful effects on an

organism. It is a function of concentration of the toxicant and the length of exposure of the animal [11]. Fish exposed to lethal doses of the xenobiotics showed drastic changes in the behavior. Respiratory upset, insensitivity to external stimuli, erratic and jerky movements along with a tendency to jump outside the convulsion were the general responses in all the treatments.

The toxic effects may include both lethal and sublethal concentration which may change the growth rate, development, reproduction, histopathology, biochemistry, physiology and behaviour [12]. The determined LC50 at 96h values of HgCl₂ was 1.85 ppm (Fig. 1). The point of Intersection with the calculated 50% mortality was determined as the TLM value Fig. 1. In present investigation, it has been endeavored to evaluate the environmental risk of man made chemicals mostly focusing on Fish due to their infusion of trial and error methods applying acute and sub lethal exposures of HgCl₂ may be explained unsuitable stress. Further, metal pollution may make mercury shift in the community structure to metal tolerant organisms, which are often capable of accumulating large amounts of metals [13].

The changes in behavior might be due to hyperactivity of the neural mechanism, as well as, for hypersecretion of stress hormones secretory endocrine glands. Studies on the median tolerance limits usually provide measures of relative toxicity of different chemicals and also the sensitivity of fish species, but not throw right on the maximum concentration to toxic materials which would not affect the metabolism of the fish. Tolerance of an organism is often measured in terms and of the level of lethal factor and length of time the animal survives.

Figure 1: Acute Toxicity Test showing tolerance of *Clarias batrachus* to Mercury

The relationship between the logarithm of the water concentrations of the test toxicant and the logarithm of fish mortality [8] appear to be linear for all the treatments and accordingly the regression lines have been drawn detecting the LC50 values (Fig. 1) by probit analysis method (Table 1). The toxicity tolerance values may vary in different size, weight and age groups of the organism in a stable environmental condition. Metal ions and their complex exhibit widening toxicity to the organism that ranges from sublethal to lethal depending upon the time of exposure and the prevailing conditions in the ambient water [14]. LC50 values differ from species to species for the same toxicant due to the mode of action and responses of the animals [15]. The toxicity tests have also been influenced by the size, age [16], sex [17] and the nutrient supply [18],[19],[20]. The present study reveals that of the two toxicants under investigation the HgCl₂ has caused great harm to the fish. This is evidenced by the facet that a slight higher concentration of HgCl₂ in the medium causes death of a much higher percentage of fish.

The lack of proportionality associated with the xenobiotics is more pronounced in mercury. Possibly, Synthesis of the detoxifying proteins (Metallothionein etc) due to cadmium induction to combat its poisoning is seemed to

be much more prominent than HgCl₂ as it also has a specific detoxification system guided by the proteins like metallothionein or glutathione. On the other hand, the inhibitory effect of HgCl₂ prevailing on the enzyme system (controller or bioaccumulation and tissue metabolism of the xenobiotics) probably gets much more prominent than that of cadmium induction.

References

- [1] Barak, 1955. Concentration of Zinc and Copper in Some Fauna from Basrah, Iraq. Mar. Res., 4: 1-6.
- [2] Chaoua, L.,1994. Studies on heavy metals in Commercial fishes from the northern part South China Sea. J. Fish. Sci. China Zhongguoshuichan – Kexue., 1: 68-74 .
- [3] Ambrose, T., Vincent, S., 1994. Susceptibility of the Fish water Fish *Gambusia affinis* (Baird and Girard), *Sarotherodon mossambicus* (Peters) and *Cirrhinus mrigala* (Ham) to Zinc toxicity. Ind. J. Environ Toxicol., 4:29-31.
- [4] APHA (2000). Standard methods for examination of water and waste water. 21st edition. American Public Health Association, Washington DC, USA. 1268.
- [5] Haniffa M.A., Porchellvi., 1984. Effect of distillery effluent oxygen consumption of fresh

- water fish *Sarotherodon mossambicus*. J. Environ. Biol., 6:17-23.
- [6] Dean, J.G., Bosqui, F.L., Lannovette, V.H., 1972. Removing heavy metals from waste water. Environ. Sci. Technol., 6: 518 – 522.
- [7] Miller, L. C., Tainter, 1944. Estimation of ED-50 its error by means of logarithmic probit graph paper. Proc. Soc. Exp. Biol. Med., 57: 261 - 264.
- [8] Finney, D.J., 1971. Probit analysis, 3rd Edn. Cambridge University Press UK , p. 25-26.
- [9] Gill, R.K., 1999. Screening of different bio materials for the removal of nickel from the industrial effluent. Ind. J. Environ. Eco plan., 2: 221 – 224.
- [10] Doudoroff, P., 1951. Doudoroff, Cherman, B.G. Anderson, G.E. Burdick, P.S. Galtsoff, Bio assay methods for the evaluation of acute toxicity of industrial wastes to fish. Sew. Ind. Wastes., 23: 1380-1397.
- [11] Alatorre-Jacome O, Garcia-Trejo F Soto-Zarazua G and E Rico-Garcia (2012). Techniques to assess the fish productivity in aquaculture farms and small fisheries: An overview of algebraic methods. J. Appl. Sci., 12(9): 888-892.
- [12] Preston, S., Coad, N., Townend, J., Killham, K., Paton, G.I., 2000. Biosensing the acute toxicity of metal interactions: are they additive, synergistic, or antagonistic? Environ. Toxicol. Chem., 19: 775-780.
- [13] Dallinger, R., Prosi, F., Segner, H., Back, H., 1987. Contaminated food uptake of heavy metals by fish: a review and a proposal for further research. Oecologia., 67: 82-89 .
- [14] Goel, P.K., 1997. Water pollution causes effects and control. New age International (P) Ltd. Publishers, New Delhi.
- [15] King, S.G., 1992. Some effects of DDT on the guppy and brown trout. Spec. Scient Res. U.S. Fish Wild. Surv., 399:1-22.
- [16] Joshi PS (2017), Dietary garlic induced growth performance, whole body composition and survival in *Clarias batrachus* (Linn.). NCRA OBS-2017:36-37
- [17] Victoriamma, D.C., Radhakrishnan., 1982. Mercury tolerance of fresh water field crab, *Ozinokeiphces senex*. Symp. Physiol. Rep. Anim. Poiluf., 67: 82-89 .
- [18] Arunachalam, S., 1980. Toxic and sublethal effects of carbaryl on the freshwater catfish, *Mystus vittatus* (Bloch). Arch. Environ. Confam. Toxicol., 9: 307-316.
- [19] Eisler, K., 1970. Factors affecting pesticide induced toxicity in estuarine fish. *Technical papers of the Bureau of sports for fish and wildlife*.
- [20] Trivedy, R.C., Dubey, P.S., 1978. Evaluation of toxicity of some industrial waste to fish by bioassay. *Environ. Pollut*, 17:75-80.